

UG/3rd Sem (H)/22/(CBCS)

2022

MATHEMATICS (Honours)

Paper Code : MTMH DC-5

(Real Analysis-II)

Full Marks : 32

Time : Two Hours

*The figures in the margin indicate full marks.
Candidates are required to give their answers
in their own words as far as practicable.*

Group - A

(4 Marks)

1. Answer any *four* questions :

1×4=4

(a) Is the function given by

$$f(x) = \begin{cases} x^2 \cos \frac{1}{2} & \text{if } 0 < x \leq 1 \\ 0 & \text{if } x = 0 \end{cases}$$

a function of bounded variation? Justify your answer.

(b) Define Refinement of a partition.

(c) Prove that $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$.

(2)

(d) If $f: [0, 1] \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} f(x) &= 0, \quad x \in [0, 1] \cap \mathbb{Q} \\ &= 1, \quad x \notin [0, 1] \cap \mathbb{Q} \end{aligned}$$

then show that f is not R-integrable on $[0, 1]$.

(e) Show that the improper integral $\int_0^1 \frac{dx}{1-x}$ is divergent.

(f) Let $f_n(x) = xe^{-nx}$, $x \geq 0$. Show that the sequence of function $\{f_n\}$ is point wise convergent on $[0, \infty)$ to the function f defined by $f(x) = 0$, $x \geq 0$.

(g) Explain why the Fundamental theorem of integral calculus can not be used to evaluate $\int_0^3 x[x]dx$.

Group - B

Answer any two questions : $5 \times 2 = 10$

2. Prove that $\lim_{x \rightarrow 0} \sum_{k=2}^{\infty} \frac{\cos kx}{k(k+1)} = \frac{1}{2}$

3. Let $f_n(x) = \log(n^2 + x^2)$, $x \in \mathbb{R}$. Show that the sequence $\{f'_n\}$ is uniformly convergent on \mathbb{R} but the sequence $\{f_n\}$ is not uniformly convergent on \mathbb{R} .

4. Prove that $f: [a, b] \rightarrow \mathbb{R}$ be a function of bounded variation on $[a, b]$ iff f can be expressed as the difference of two monotonic increasing functions on $[a, b]$.

5. Let $f: [a, b] \rightarrow \mathbb{R}$ be integrable on $[a, b]$. If there exists a positive real number K such that $f(x) \geq K$ for all $x \in [a, b]$ then show that $\frac{1}{f}$ is integrable on $[a, b]$.

Group - C

Answer any two questions : $9 \times 2 = 18$

6. (a) Let $f: [a, b] \rightarrow \mathbb{R}$ be bounded on $[a, b]$ and let f be continuous on $[a, b]$ except on a infinite subset $S \subset [a, b]$ such that the number of limit points of S is finite. Then prove that f is R -integrable on $[a, b]$. 4

- (b) Prove that the even function $f(x) = |x|$ on $[-\pi, \pi]$ has as cosine series in Fourier's form as

$$\frac{\pi}{2} - \frac{4}{\pi} \left\{ \cos x + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right\}. \quad 5$$

7. (a) If $f: [a, b] \rightarrow \mathbb{R}$ be integrable on $[a, b]$ and f possesses an antiderivative ϕ on $[a, b]$, then prove that $\int_a^b f = \phi(b) - \phi(a)$

[fundamental theorem of Integral Calculus]. 5

- (b) Find the length of the perimeter of the cardioid $r = a(1 + \cos \theta)$. 4

8. (a) Test the convergence of β function. 5

- (b) State and prove the Cauchy-Hadamard theorem. 4

UG/3rd Sem (H)/22/(CBCS)

2022

MATHEMATICS (Honours)

Paper Code : MTMH DC-6

(Linear Algebra)

Full Marks : 32

Time : Two Hours

The figures in the margin indicate full marks.

*Candidates are required to give their answers
in their own words as far as practicable.*

Group - A

(4 Marks)

1. Answer any *four* questions : 1×4=4

(a) Find $k \in \mathbb{R}$ so that the set $S = \{(1, 2, 1), (k, 3, 1), (2, k, 0)\}$ is linearly dependent in \mathbb{R}^3 .

(b) Find the dimension of the subspace S of \mathbb{R}^4 defined by

$$S = \{(x, y, z, w) \in \mathbb{R}^4 : x + y + z + w = 0\}$$

(c) If α, β be two orthogonal vectors in a Euclidean space V , then show that

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2$$

P.T.O.

(2)

(d) Find the range of the linear transformation

$T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ given by

$$T(x, y, z) = (x + z, x + y + 2z, 2x + y + 3z)$$

(e) If $\{u_1, u_2, \dots, u_r\}$ be an orthonormal set, prove that for any $v \in V$, the vector $w = v - \langle v, u_1 \rangle u_1 - \langle v, u_2 \rangle u_2 - \dots - \langle v, u_r \rangle u_r$ is orthogonal to each of the u_i .

(f) Let λ be an eigenvalue of a linear operator T on an inner product space V . If $T^* = T^{-1}$, then show that $|\lambda| = 1$.

(g) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation given by

$$T(x_1, x_2) = (x_1 + x_2, x_1 - x_2, x_2)$$

Then find rank T .

Group - B

(10 Marks)

Answer any two questions.

5×2=10

2. If U and W be two subspaces of a vector space V over a field F such that $U \cap W = \{\theta\}$ and if $\{\alpha_1, \alpha_2, \dots, \alpha_m\}$ and $\{\beta_1, \beta_2, \dots, \beta_n\}$ be respectively the bases of U and W , then show that $\{\alpha_1, \alpha_2, \dots, \alpha_m, \beta_1, \beta_2, \dots, \beta_n\}$ is a basis of $U + W$. 5

(3)

3. Determine the linear mapping $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ which maps the basis vectors $(0, 1, 1)$, $(1, 0, 1)$, $(1, 1, 0)$ of \mathbb{R}^3 to $(1, 1, 1)$, $(1, 1, 1)$, $(1, 1, 1)$ respectively. Verify that $\dim \ker T + \dim \text{Im} T = 3$. 5

4. Find the algebraic and geometric multiplicities of each eigen value of the matrix

$$\begin{pmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{pmatrix} \quad 5$$

5. If $T: V \rightarrow V$ be a linear transformation, show that the following statements are equivalent :

(i) $\text{Range } T \cap \text{Ker} T = \{0\}$

(ii) If $T(T(v)) = 0$ then $T(v) = 0, v \in V$. 3+2

Group - C

(18 Marks)

Answer any two questions. 9×2=18

6. (a) If V is a finite dimensional inner product space and W is a subspace of V , then show that $V = W \oplus W^\perp$. 5

(b) Let T be a normal operator. Prove :

(i) $T(v) = 0$ if and only if $T^*(v) = 0$

(ii) $T - \lambda I$ is normal. 2+2

P.T.O.

7. (a) Extend the set of vectors $\{(2, 3, -1), (1, -2, -4)\}$ to an orthogonal basis of the Euclidean space \mathbb{R}^3 with standard inner product and then find the associated orthonormal basis. 5

(b) Let T be the linear operator on \mathbb{R}^3 defined by $T(x_1, x_2, x_3) = (3x_1 + x_3, -2x_1 + x_2, -x_1 + 2x_2 + 4x_3)$. Show that T is invertible. 4

8. (a) Apply gram-schmidt process to obtain an orthonormal basis of the subspace of the Euclidean space \mathbb{R}^4 with standard inner product, spanned by the vectors $(1, 1, 0, 1), (1, -2, 0, 0), (1, 0, -1, 2)$. 5

(b) A linear mapping $T: \mathbb{R}^3 \rightarrow \mathbb{R}^4$ is defined by $T(x_1, x_2, x_3) = (x_2 + x_3, x_3 + x_1, x_1 + x_2, x_1 + x_2 + x_3)$, $(x_1, x_2, x_3) \in \mathbb{R}^3$. Find $\ker T$. Verify that the set $\{T(\epsilon_1), T(\epsilon_2), T(\epsilon_3)\}$ is linearly independent in \mathbb{R}^4 , where $\epsilon_1 = (1, 0, 0)$, $\epsilon_2 = (0, 1, 0)$ and $\epsilon_3 = (0, 0, 1)$. 4

UG/3rd Sem (H)/22/(CBCS)

2022

MATHEMATICS (Honours)

Paper Code : MTMH DC-7

(Multivariate Calculus & Vector Calculus)

Full Marks : 32

Time : Two Hours

*The figures in the margin indicate full marks.
Candidates are required to give their answers
in their own words as far as practicable.*

Symbols used in this question paper bears
their original meaning unless stated.

Group - A

(4 Marks)

1. Answer any *four* questions :

1×4=4

(a) Show that $\lim_{\substack{x \rightarrow 0 \\ y \rightarrow 0}} f(x, y)$ does not exist, where

$$f(x, y) = \begin{cases} \frac{x^2 - y^2}{x^2 + y^2}, & x^2 + y^2 \neq 0 \\ 0, & x^2 + y^2 = 0 \end{cases}$$

(b) If $f(x, y) = \begin{cases} \frac{x^3 - y^3}{x^2 + y^2}, & x^2 + y^2 \neq 0 \\ 0, & x^2 + y^2 = 0, \end{cases}$

then show that $f_x(0, 0) \neq f_y(0, 0)$.

P.T.O.

(2)

(c) Find the directional derivative of

$$\varphi = 4xz^3 - 3x^2y^2z \text{ at } (2, -1, 2) \text{ in the direction } 2\hat{i} - 3\hat{j} + 6\hat{k}.$$

(d) Evaluate $\int_0^{\frac{\pi}{2}} \int_0^{\cos\theta} drd\theta$.

(e) Evaluate $\iiint_V dx dy dz$ where V is the tetrahedron bounded by $x = 0, y = 0, z = 0, x + y + z = 1$.

(f) Prove that $\text{Curl}(\text{grad } \varphi) = \vec{0}$

(g) State Gauss divergence theorem.

Group - B

(10 Marks)

Answer any *two* questions : $5 \times 2 = 10$

2. Show that the necessary and sufficient condition that a nonzero differentiable vector function $\vec{f}(t)$ to possess the constant magnitude is that $\vec{f} \cdot \frac{d\vec{f}}{dt} = 0$.

3. Evaluate the line integral $\oint_{\Gamma} \vec{f} \cdot d\vec{r}$ by Stokes theorem where Γ being the boundary of the rectangle $0 \leq x \leq \frac{\pi}{2}, 0 \leq y \leq 2, z = 1$ and $\vec{f} = \sin z\hat{i} - \cos x\hat{j} + \sin y\hat{k}$.

(3)

4. Show that $f(x, y) = \begin{cases} \frac{x^2 y}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0 & , (x, y) = (0, 0) \end{cases}$

has directional derivative at $(0, 0)$ in any direction $\beta = (l, m)$ where $l^2 + m^2 = 1$ but f is discontinuous at $(0, 0)$.

5. Show that if $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$, the maximum value of xyz is $\frac{abc}{3\sqrt{3}}$.

Group - C

(18 Marks)

Answer any two questions : 9×2=18

6. (a) State and prove Schwarts Theorem. 6

(b) Show that $\int_0^1 \int_0^{1-y^2} \left\{ (x-1)^2 + y^2 \right\} dx dy = \frac{44}{105}$. 3

7. (a) Show that $\bar{\nabla} \log r = \frac{1}{r^2} \bar{r}$

where $\bar{r} = x\hat{i} + y\hat{j} + z\hat{k}$ and $r = |\bar{r}|$. 4

(b) Use Greens theorem to evaluate $\oint_{\Gamma} (xy dx + y^2 dy)$ where Γ is a square in the xy plane with vertices $(0, 0)$, $(0, 1)$, $(1, 0)$ and $(1, 1)$. 5

P.T.O.

(4)

8. (a) Let $V = \sin^{-1} \sqrt{\frac{x^{\frac{1}{3}} + y^{\frac{1}{3}}}{x^{\frac{1}{2}} + y^{\frac{1}{2}}}}$. Prove that

$$x^2 \frac{\partial^2 V}{\partial x^2} + 2xy \frac{\partial^2 V}{\partial x \partial y} + y^2 \frac{\partial^2 V}{\partial y^2} = \frac{\tan V}{12} \left(\frac{13}{12} + \frac{\tan^2 V}{12} \right).$$

5

(b) If $\vec{f} = t^2 \hat{i} - t \hat{j} + (2t+1) \hat{k}$ and $\vec{g} = (2t-3) \hat{i} + \hat{j} - t \hat{k}$

then prove that $\frac{d}{dt} \left(\vec{f} \times \frac{d\vec{g}}{dt} \right) = \hat{i} + 6\hat{j} + 2\hat{k}$ at

$t = 1$.

4